arXiv:1404.4211v1 [math.NT] 16 Apr 2014

IMAGES OF POLYNOMIAL MAPS ON LARGE FIELDS

MICHIEL KOSTERS

Abstract. A field k is called large if every irreducible k-curve with a k-rational smooth point has infinitely many k-points. Let k be a perfect large field and let $f \in k[x]$. Consider the evaluation map $f_k : k \to k$. Assume that f_k is not surjective. We will show that $k \setminus f_k(k)$ is infinite. This conclusion follows from a similar statement about finite morphisms between normal projective curves over perfect large fields.

1. Introduction

Problem 1. Let k be an infinite field and let $f \in k[x]$. Consider the evaluation map $f_k : k \to k$. Assume that f_k is not surjective. Is $k \setminus f_k(k)$ infinite?

This question was asked by Philipp Lampe on mathoverflow as Question 6820 in 2009. The problem is still unsolved in general. It is not hard to see that the problem has a positive answer when k is Hilbertian. Number fields are for example Hilbertian. In this article we will positively answer this problem for perfect large fields. Let us first define what a large field is.

Definition 1.1. A field k is called *large* if every irreducible k-curve C with a k-rational smooth point has infinitely many k-points.

Note that large fields are infinite. For more information on large fields see the survey [Pop13]. Some examples of large fields are $\mathbf{R}$, $\mathbf{Q}_p$ (p prime), $l((t))$ (where l is a field), infinite algebraic extensions of finite fields. Furthermore, finite extensions of large fields are large. In the definition of large field we can replace the word smooth by normal if k is perfect ([Liu02, Corollary 4.3.33 and Lemma 8.2.21])

Our main theorem is the following (we follow [Liu02] for the definition of a normal projective curve).

Theorem 1.2. *Let k be a perfect large field. Let C, D be normal projective curves over k. Let $f\colon C \to D$ be a finite morphism. Suppose that the induced map $f_k\colon C(k) \to D(k)$ is not surjective. Then one has $|D(k) \setminus f_k(C(k))| = |k|$.*

The following corollary gives a positive answer to our problem for perfect large fields, since large fields are infinite.

Corollary 1.3. *Let k be a perfect large field. Then the following hold.*

i. *Let $f \in k(x)$ such that the induced map $f_k\colon \mathbf{P}^1_k(k) \to \mathbf{P}^1_k(k)$ is not surjective. Then one has $|\mathbf{P}^1_k(k) \setminus f_k(\mathbf{P}^1_k(k))| = |k|$.*
ii. *Let $f \in k[x]$ such that the induced evaluation map $f_k\colon k \to k$ is not surjective. Then one has $|k \setminus f_k(k)| = |k|$.*

Date: April 17, 2014.
2010 *Mathematics Subject Classification.* 12-02, 12J10, 11R58.
This article is part of my PhD thesis written under the supervision of Hendrik Lenstra.

2. Prerequisites

2.1. **Valuations and curves.**

Definition 2.1. Let K be a field. Then a *valuation ring* on K is a subring $\mathcal{O}$ of K such that for all $x \in K^*$ one has $x \in \mathcal{O}$ or $x^{-1} \in \mathcal{O}$.

Such a valuation ring is a local ring and it is integrally closed in K. For notational convenience, we often say that P is a valuation with corresponding valuation ring $\mathcal{O}_P$, maximal ideal $\mathfrak{m}_P$ and residue field k_P. We say that (K, P) is a valued field.

Let (K, P) be a valued field. Let L be a field extension of K. A valuation Q on L is an extension of P if $\mathcal{O}_Q \cap K = \mathcal{O}_P$. We denote such an extension by $(K, P) \subseteq (L, Q)$. Such an extension is called finite (Galois, normal, ...) if L/K is finite (Galois, normal, ...).

Definition 2.2. Let k be a field. A *function field* over k is a finitely generated field extension of k of transcendence degree 1.

Theorem 2.3. *There is an anti-equivalence of categories between the category of normal projective curves over k with finite morphisms and the category of function fields over k with finite k-morphisms of fields. This equivalence maps a curve C to its function field $k(C)$ and a finite morphism $C \to D$ to the induced inclusion $k(D) \subseteq k(C)$.*

Proof. See [Liu02, Proposition 7.3.13]. □

Let C be a normal projective curve over a field k. Each closed point P gives rise to a valuation on $k(C)$ which contains k but is not equal to $k(C)$, namely $\mathcal{O}_P$. Conversely, every such valuation corresponds to a point. This set of such valuations is denoted by $\mathcal{P}_{K/k}$. We let $\mathcal{P}^1_{K/k} \subseteq \mathcal{P}_{K/k}$ be the subset consisting of valuations P which satisfy $\mathrm{k}_P = k$. When $K = k(C)$, the set $\mathcal{P}^1_{K/k}$ corresponds to $C(k)$.

2.2. **Galois theory of valuations.**

We refer to [Kos14] for a Galois theoretic approach to valuation theory. We recall some of the important statements.

Let K be a field with valuation P and let M be a normal extension of K with group $G = \mathrm{Aut}_K(M)$. Let Q be an extension of P to M. Set $\mathrm{D}_{Q,K} = \{g \in G : gQ = Q\}$ (*decomposition group*). Note that we have a natural map $\mathrm{D}_{Q,K} \to \mathrm{Aut}_{\mathrm{k}_P}(\mathrm{k}_Q)$. The kernel of this map is called the *inertia group* and is denoted by $\mathrm{I}_{Q,K}$.

The following theorem summarizes the results from general valuation theory which we need later on.

Theorem 2.4. *Let L be an intermediate extension of M/K. The following hold:*

i. *The set of valuations extending P to M is not empty and the group G acts transitively on it.*
ii. *The sequence*
$$0 \to \mathrm{I}_{Q,K} \to \mathrm{D}_{Q,K} \to \mathrm{Aut}_{\mathrm{k}_P}(\mathrm{k}_Q) \to 0$$
is exact. It is right split (as topological groups) if k_Q is algebraically closed.
iii. *One has $\mathrm{D}_{Q,L} = \mathrm{Aut}_L(M) \cap \mathrm{D}_{Q,K}$ and $\mathrm{I}_{Q,L} = \mathrm{Aut}_L(M) \cap \mathrm{I}_{Q,K}$.*
iv. *If L/K is Galois, then then the maps $\mathrm{D}_{Q,K} \to \mathrm{D}_{Q|_L,K}$ and $\mathrm{I}_{Q,K} \to \mathrm{I}_{Q|_L,K}$ are surjective.*

Proof. i: See [Kos14, Proposition 5.6].
ii: See [Kos14, Theorem 3.8iii].
iii: Follows directly from the definitions.
iv: See [Kos14, Theorem 3.6]. □

One of the reasons for assuming that k is perfect in our main theorem in the following proposition.

Proposition 2.5. *Let (K, P) be a valued field and let L be a finite algebraic extension of K. Assume that k_P is perfect. Let $(M, Q) \supseteq (K, P)$ be a finite normal extension of valued fields with group $G = \mathrm{Aut}_K(M)$ such that the G-set $X = \mathrm{Hom}_K(L, M)$ is not empty. Then the cardinality of the set of valuations P' on L extending P such that $\mathrm{k}_{P'} = \mathrm{k}_P$ is equal to $\#\left(\mathrm{I}_{Q,K} \backslash X\right)^{\mathrm{D}_{Q,K} \,/\, \mathrm{I}_{Q,K}}$.*

Proof. See [Kos14, Corollary 3.18]. □

2.3. **Function fields and ramification.**

Proposition 2.6. *Let K be a function field over a perfect field k. Let $\overline{k}$ be an algebraic closure of k in an algebraic closure of K. The following hold:*

i. *Let M/K be a finite normal extension. Then only finitely many primes $P \in \mathcal{P}_{K/k}$ have an extension $Q \in \mathcal{P}_{M/k}$ with the property that $\mathrm{I}_{Q,K} \neq 0$.*
ii. *For all primes $Q \in \mathcal{P}_{\overline{k}K/k}$ one has $\mathrm{I}_{Q,K} = 0$. Furthermore, if k is integrally closed in K, then a prime $P \in \mathcal{P}_{K/k}$ has a unique extension to $\mathcal{P}_{\overline{k}K/k}$ if and only if $P \in \mathcal{P}^1_{K/k}$.*

Proof. i. Since valuation extend unique in purely inseparable field extensions, we may assume that M/K is Galois. In that case, the statement is well-known: only finitely many primes ramify.
ii. This follows from [Sti09, Theorem 3.6.3].

□

3. Proofs of the theorems

Definition 3.1. Let K be a function field over a field k and let $P \in \mathcal{P}^1_{K/k}$. Let $\overline{K}$ be an algebraic closure of K. Let M/K be a normal extension of K inside $\overline{K}$ with group $G = \mathrm{Aut}_K(M)$. Let $\overline{k}$ be the algebraic closure of k inside $\overline{K}$. Set $\Gamma = \mathrm{Aut}_k(\overline{k})$. Let Q be an extension of P to M. A *Frobenius* for Q/P is a continuous morphism $\Gamma \to G$ such that that the diagram, where ψ is the natural map,

$$\begin{array}{ccc} \Gamma & \xrightarrow{\quad\varphi\quad} & G \\ & \searrow^{\psi} & \downarrow{\scriptstyle \pi} \\ & & \mathrm{Aut}(\overline{k} \cap M/k) \cong \mathrm{Aut}((M \cap (\overline{k}K))/K) = G/N \end{array}$$

commutes and such that $\mathrm{im}(\varphi)\, \mathrm{I}_{Q,K} = \mathrm{D}_{Q,K}$. The set of all such Frobenius maps is denoted by $\mathrm{Frob}(Q/P)$.

For the next proposition and lemma assume that M/K is a normal extension of function fields over a perfect field k inside $\overline{K}$. Let $G = \mathrm{Aut}_K(M)$. Let $P \in \mathcal{P}^1_{K/k}$ with Q above it in M.

Suppose $\varphi \in \mathrm{Frob}(Q/P)$. We use the notation as in the definition of a Frobenius. Consider $\mathrm{Graph}(\varphi) \subseteq \Gamma \times_{G/N} G = \mathrm{Gal}(\overline{k}M/K)$, which is a closed subgroup. Set $M^\varphi = (\overline{k}M)^{\mathrm{Graph}(\varphi)}$. Note that M^φ/K is a finite extension if M/K is finite. Furthermore, M^φ is geometrically irreducible over k since k is perfect. Finally, we have $\overline{k}M^\varphi = \overline{k}M$.

Proposition 3.2. *There is $\varphi \in \mathrm{Frob}(Q/P)$ such that $\mathcal{P}^1_{M^\varphi/k}$ is not empty.*

Proof. Let $\overline{Q}$ be an extension of Q to $\overline{k}M$. We have the following split exact sequence (Theorem 2.4ii):

$$0 \to \mathrm{I}_{\overline{Q},K} \to \mathrm{D}_{\overline{Q},K} \to \Gamma \to 0.$$

Let $\varphi_0 \colon \Gamma \to \mathrm{D}_{\overline{Q},K} \subseteq \Gamma \times_{G/N} G$ be such a splitting. Note that $\mathrm{im}(\varphi_0)$ is the graph of a function $\varphi \colon \Gamma \to \mathrm{D}_{Q,K}$. One has $\mathrm{im}(\varphi_0)\,\mathrm{I}_{\overline{Q},K} = \mathrm{D}_{\overline{Q},K}$. This gives $\mathrm{im}(\varphi)\,\mathrm{I}_{Q,K} = \mathrm{D}_{Q,K}$ as required (Theorem 2.4iv). The commutativity of the diagram follows since $\mathrm{Graph}(\varphi) = \mathrm{im}(\varphi_0) \subseteq \Gamma \times_{G/N} G$.

We will show $Q' = \overline{Q}|_{M^\varphi} \in \mathcal{P}^1_{M^\varphi/k}$. By construction we have $\mathrm{D}_{\overline{Q},K} \supseteq \mathrm{Graph}(\varphi)$. Hence we obtain $\mathrm{D}_{\overline{Q},M^\varphi} = \mathrm{D}_{\overline{Q},K} \cap \mathrm{Gal}(M/M^\varphi) = \mathrm{Gal}(M/M^\varphi)$ (Theorem 2.4iii). Hence $\overline{Q}$ is the unique prime above Q' in $\overline{k}M$ (Theorem 2.4i) and Q' is rational (Proposition 2.6ii). □

Lemma 3.3. *Let $\varphi \in \mathrm{Frob}(Q/P)$. Consider the natural map $\psi \colon \mathcal{P}^1_{M^\varphi/k} \to \mathcal{P}^1_{K/k}$. Let $P \in \mathrm{im}(\psi)$ with the property that any extension Q of P to M satisfies $\mathrm{I}_{Q,K} = 0$. Then there is a prime Q of M above P with $\mathrm{D}_{Q,K} = \mathrm{im}(\varphi)$.*

Proof. Let Q'' be a valuation on $\overline{k}M$ extending P such that $Q' = Q''|_{M^\varphi} \in \mathcal{P}^1_{M^\varphi/k}$ (Theorem 2.4i). Note that $\mathrm{I}_{Q'',K} = 0$ (this follows from Proposition 2.6). Set $Q = Q''|_M$ and $P' = Q''_{\overline{k}K}$. As Q' is rational, the natural injective map

$$\mathrm{D}_{Q'',K} \cap \mathrm{Graph}(\varphi) = \mathrm{D}_{Q'',M^\varphi} \cong \mathrm{Aut}(k_{Q''}/k_{Q'}) \to \mathrm{D}_{Q'',K} \cong \mathrm{Aut}(k_{Q''}/k)$$

is surjective (Theorem 2.4iv). Hence we find $\mathrm{D}_{Q'',K} \subseteq \mathrm{Graph}(\varphi)$. The map $\mathrm{D}_{Q'',K} \to \mathrm{D}_{P',K} = \Gamma$ is surjective (Theorem 2.4iv, [Sti09, Theorem 3.6.3]). As $\mathrm{Graph}(\varphi)$ is a graph, this shows that $\mathrm{D}_{Q'',K} = \mathrm{Graph}(\varphi)$. We deduce $\mathrm{D}_{Q,K} = \mathrm{im}(\varphi)$ (Theorem 2.4iv). □

Proof of Theorem 1.2. With the help of Theorem 2.3 we see that equivalently we need to prove the following. Let L/K be a finite extension of function fields over k. Assume that the induced map $f_k \colon \mathcal{P}^1_{L/k} \to \mathcal{P}^1_{K/k}$ is not surjective. Show that $|\mathcal{P}^1_{K/k} \setminus \mathrm{im}(f_k)| = |k|$.

Let M be a finite normal extension of K such that $X = \mathrm{Hom}_K(L, M) \neq \emptyset$. Assume $P \in \mathcal{P}^1_{K/k}$, $P \notin \mathrm{im}(f_k)$. Let Q be an extension of P to M. Let $\varphi \in \mathrm{Frob}(Q/P)$ with $\mathcal{P}^1_{M^\varphi/k} \neq \emptyset$ (Proposition 3.2). Since k is a large field, one has $|\mathcal{P}^1_{M^\varphi/k}| = |k|$ ([Jar11, Proposition 5.4.3]). Note that we have $\mathrm{im}(\varphi)\,\mathrm{I}_{Q,K} = \mathrm{D}_{Q,K}$. As $P \notin \mathrm{im}(f_k)$ we conclude from Corollary 2.5 that $(\mathrm{I}_{Q,K} \setminus X)^{\mathrm{D}_{Q,K}\,/\,\mathrm{I}_{Q,K}} = \emptyset$. This implies $X^{\mathrm{im}(\varphi)} = \emptyset$. Consider the map $\psi \colon \mathcal{P}^1_{M^\varphi/k} \to \mathcal{P}^1_{K/k}$. Let T be the set of $P' \in \mathrm{im}(\psi)$ such that for any extension Q' of P' to M one has $\mathrm{I}_{Q',K} = 0$. Then T has cardinality $|k|$ as well (Propostion 2.6i, Theorem 2.4i). For $P' \in T$ there is a

prime Q' of M above P' with $\mathrm{im}(\varphi) = \mathrm{D}_{Q',K}$ (Lemma 3.3). We find

$$(\mathrm{I}_{Q',K} \setminus X)^{\mathrm{D}_{Q',K} \,/\, \mathrm{I}_{Q',K}} = X^{\mathrm{D}_{Q',K}} = X^{\mathrm{im}(\varphi)} = \emptyset.$$

From Corollary 2.5 we conclude that $P' \notin \mathrm{im}(f_k)$. This finishes the proof. □

Proof of Corollary 1.3. i: The statement is true if f is constant ([Jar11, Proposition 5.4.3]). Assume $f \in k(x)$ is not constant. Then f induces a finite morphism $\mathbf{P}^1_k \to \mathbf{P}^1_k$ and we apply Theorem 1.2.

ii: This follows from i, since the map sends the point at infinity to the point at infinity. □

References

[Jar11] Moshe Jarden. *Algebraic patching*. Springer Monographs in Mathematics. Springer, Heidelberg, 2011.

[Kos14] Michiel Kosters. The algebraic theory of valued fields. `http://arxiv.org/abs/1404.3916`, 2014. preprint.

[Liu02] Qing Liu. *Algebraic geometry and arithmetic curves*, volume 6 of *Oxford Graduate Texts in Mathematics*. Oxford University Press, Oxford, 2002. Translated from the French by Reinie Erné, Oxford Science Publications.

[Pop13] Florian Pop. Little survey on large fields. 2013.

[Sti09] Henning Stichtenoth. *Algebraic function fields and codes*, volume 254 of *Graduate Texts in Mathematics*. Springer-Verlag, Berlin, second edition, 2009.

Mathematisch Instituut P.O. Box 9512 2300 RA Leiden the Netherlands

E-mail address: `mkosters@math.leidenuniv.nl`

URL: `www.math.leidenuniv.nl/~mkosters`